Summary of Results

PATIENT NAME: Last, First

DOB: **23-Jan-1922**



GENDER: Female
SPECIMEN ID: SID 234/6-A
PATIENT/MRN: MRN 123456
CUSTOMER REF: CR 98765

ORDERED BY: M.D. Irvine - IT Test, Doctor ACCOUNT: IRV - Agendia IT Test

12345 Main St Some City CA

99999 US

REQUISITION #: R# 2345678
SPECIMEN TYPE: FFPE, Needle Core
SPECIMEN SOURCE: Right Medial Breast
COLLECTED DATE: 13-Jan-2015
RECEIVED DATE: 14-Jan-2015
REPORTED DATE: 27-Jan-2015


Summary of Results: High Risk HER2-type


Risk of Recurrence

High Risk

Molecular Subtype

HER2-type





Note: This information is provided for general informational purposes. It is not part of any official diagnostic report. Please refer to individual MammaPrint and BluePrint reports for comments, assay information, disclaimer and references.

ustomercare@agendia.com | www.agendia.com | © 2015 FFP16-000288 M-USA-001-V3



PATIENT NAME: Last, First

REPORTED DATE: 27-Jan-2015

p<0.01

Distant Metastasis Free Survival @ 5 yrs for MammaPrint High Risk² Endocrine + Chemotherapy Endocrine 50% 55% 60% 65% 70% 75% 80% 85% 90% 95% 100%


• The MammaPrint result provides independently validated, statistically significant, additive information for physicians to assist them in making treatment decisions for early stage breast cancer patients.

DMFS %

- If the risk assessment by MammaPrint and clinicopathological characteristics is concordant and indicates a High Risk of recurrence, the use of combined
 endocrine and chemotherapy (ET+CT) seems clinically indicated.
- If the risk assessment by MammaPrint and clinicopathological characteristics is discordant, MammaPrint High Risk and clinically stratified Low Risk patients will likely benefit from chemotherapy. If these patients are highly endocrine-responsive (≥50% ER positivity), endocrine therapy (ET) alone might be the desired option; however, withholding chemotherapy might not be supported due to the observed improvement in DMFS at 5 years for MammaPrint High Risk patients who received ET+CT.
- Other factors, such as age and co-morbidities, may influence the decision-making process for systemic adjuvant therapy shared between the physicians and
 patients. Distant metastasis-free survival (DMFS) is defined as time from surgery to any distant metastasis.

Estimated benefit in breast cancer specific survival by trastuzumab:

For women with early-stage HER2-positive breast cancer, addition of trastuzumab to paclitaxel after doxorubicin and cyclophosphamide results in a 10-year absolute benefit of 9% in overall survival (OS) and 11% in disease-free survival (DFS).



References: (1) Buyse M, Loi S, van't Veer L et al., J Natl Cancer Inst. 2006;98(17):1183-92. (2) Knauer M, Mook S, Rutgers EJ et al., Breast Cancer Res Treat. 2010;120(3):655-61. (3) Perez EA, Romond EH, Suman VJ, et al., J Clin Oncol. 2014;32(33):3744-52. (4) Gluck S, de Snoo F, Peeters J et al., Breast Cancer Res Treat. 2013;139(3):759-67.

Agendia Summary Page

Disclaimer: The summary page is provided for general informational purposes only and is not part of any official diagnostic report. Please refer to the official individual patient reports for final results. This information (including, without limitation, advice and recommendations) and services are neither medical nor health care advice for any individual problem nor a substitute for advice and services from a qualified health care provider familiar with the patient's medical history. All publication information can be found at www.agendia.com.