

Summary of Results

PATIENT NAME: **Last Name, First Name**

DOB: **10-Jan-1961**

GENDER: Female
SPECIMEN ID: MRN 123456
PATIENT/MRN: 945839302
CUSTOMER REF: 123456789

ORDERED BY: Dr. Doe, John
ACCOUNT: John Doe Hospital
1234 Main St.
Irvine CA 92618 USA

REQUISITION #: 1234567
SPECIMEN TYPE: FFPE, Core
SPECIMEN SOURCE: Left Breast
COLLECTED DATE: 18-Feb-2014
RECEIVED DATE: 19-Feb-2014
REPORTED DATE: 21-Feb-2014

Summary of Results: **Low Risk Luminal-type (A)**


Risk of Recurrence

Low Risk


Molecular Subtype

Luminal-type

MammaPrint[®] FFPE: 70-Gene Breast Cancer Recurrence Assay



Blueprint[®]: 80-Gene Molecular Subtyping Assay




Note: This information is provided for general informational purposes. It is not part of any official diagnostic report. Please refer to individual MammaPrint and Blueprint reports for comments, assay information, disclaimer and references.

PATIENT NAME: **Jane Doe-Jane Doe-Jane**

REPORTED DATE: **21-Feb-2014**

Adjuvant Response to Therapy




- The MammaPrint result provides independently validated, statistically significant, additive information for physicians to assist them in making treatment decisions for early stage breast cancer patients.
- If the risk assessment by MammaPrint and clinicopathological characteristics is concordant and indicates a Low Risk of recurrence, endocrine therapy (ET) alone should be adequate treatment.
- If the risk assessment by MammaPrint and clinicopathological characteristics is discordant, MammaPrint Low Risk and clinically stratified High Risk patients will likely benefit from ET alone for highly endocrine-responsive patients (≥50% ER positivity), as defined by the 2009 St. Gallen consensus panel. Since the risk of recurrence for these patients is so low, they will likely gain little or no benefit from additional chemotherapy (CT).
- Other factors, such as age and co-morbidities, may influence the decision-making process for systemic adjuvant therapy shared between the physicians and patients. Distant metastasis-free survival (DMFS) is defined as time from surgery to any distant metastasis.

Estimated benefit in breast cancer specific survival by trastuzumab:

For women with early-stage HER2-positive breast cancer, addition of trastuzumab to paclitaxel after doxorubicin and cyclophosphamide results in a 10-year absolute benefit of 9% in overall survival (OS) and 11% in disease-free survival (DFS).³


Neoadjuvant Response to Therapy

Low Risk Luminal-type (A) Neoadjuvant Chemosensitivity⁴



Subtype Results	Chemosensitivity Relevance
Low Risk Luminal-type (A)	<ul style="list-style-type: none"> • Low likelihood of pCR • No expected benefit from chemotherapy • Endocrine therapy further reduces risk

Distant Metastasis-Free Survival (DMFS) by Molecular Subtype



References: (1) Buyse M, Loi S, van't Veer L et al., J Natl Cancer Inst. 2006;98(17):1183-92. (2) Knauer M, Mook S, Rutgers EJ et al., Breast Cancer Res Treat. 2010;120(3):655-61. (3) Perez EA, Romond EH, Suman VJ, et al., J Clin Oncol. 2014;32(33):3744-52. (4) Gluck S, de Snoo F, Peeters J et al., Breast Cancer Res Treat. 2013;139(3):759-67.

Agendia Summary Page

Disclaimer: The summary page is provided for general informational purposes only and is not part of any official diagnostic report. Please refer to the official individual patient reports for final results. This information (including, without limitation, advice and recommendations) and services are neither medical nor health care advice for any individual problem nor a substitute for advice and services from a qualified health care provider familiar with the patient's medical history. All publication information can be found at www.agendia.com.